arXiv:1304.4394v1 [math.CA] 16 Apr 2013

BVP's with discontinuities at finite number interior points and spectral parameter in the boundary conditions

K. Aydemir[a]

[a]*Department of Mathematics, Faculty of Arts and Science, Gaziosmanpaşa University, 60250 Tokat, Turkey*

Abstract

In this study we are concerned with a class of generalized BVP' s consisting of eigendependent boundary conditions and supplementary transmission conditions at finite number interior points. By modifying some techniques of classical Sturm-Liouville theory and suggesting own approaches we find asymptotic formulas for the eigenvalues and eigenfunction.

Keywords: Sturm-Liouville problems, eigenvalue, eigenfunction, asymptotics of eigenvalues and eigenfunction.

---

1. Introduction

In this study we shall investigate a new class of BVP's which consist of the Sturm-Liouville equation

$$\mathcal{L}(u) := -\rho(x)u''(x) + q(x)u(x) = \lambda u(x), x \in \Omega \tag{1}$$

together with eigenparameter-dependent boundary conditions at end points $x = a, b$

$$\mathcal{L}_1(u) := \alpha_1 u(a) - \alpha_2 u'(a) - \lambda(\alpha_3 u(a) - \alpha_4 u'(a)) = 0, \tag{2}$$

$$\mathcal{L}_2(u) := \beta_1 u(b) - \beta_2 u'(b) + \lambda(\beta_3 u(b) - \beta_4 u'(b)) = 0, \tag{3}$$

---

[*] Corresponding Author (Tel: +90 356 252 16 16, Fax: +90 356 252 15 85)
Email address: `kadriye.aydemir@gop.edu.tr` (K. Aydemir)

and the transmission conditions at finite interior points $\xi_i \in (a,b), i = 1,2,...r$

$$\pounds_i(u) = \alpha_{i1}^{-}u'(\xi_i-) + \alpha_{i0}^{-}u(\xi_i-) + \alpha_{i1}^{+}u'(\xi_i+) + \alpha_{i0}^{+}u(\xi_i+) = 0, \tag{4}$$

$$\mathfrak{L}_i(u) = \beta_{i1}^{-}u'(\xi_i-) + \beta a_{i0}^{-}u(\xi_i-) + \beta_{i1}^{+}u'(\xi_i+) + \beta_{i0}^{+}u(\xi_i+) = 0, \tag{5}$$

where $\Omega = \bigcup\limits_{i=1}^{r+1}(\xi_{i-1},\xi_i),\ a := \xi_0,\ b := \xi_{r+1},\ \rho(x) = \rho_i^2 > 0$ for $x \in \Omega_i := (\xi_{i-1},\xi_i),\ i = 1,2,...r+1$, the potential $q(x)$ is real-valued function which continuous in each of the intervals (ξ_{i-1},ξ_i), and has a finite limits $q(\xi_i \mp 0)$, λ is a complex spectral parameter, $\alpha_k,\ \beta_k\ (k = 1,2,3,4),\ \alpha_{ij}^{\pm},\ \beta_{ij}^{\pm}\ (i = 1,2,...r$ and $j = 0,1)$ are real numbers. We want emphasize that the boundary value problem studied here differs from the standard boundary value problems in that it contains transmission conditions and the eigenvalue-parameter appears not only in the differential equation, but also in the boundary conditions. Moreover the coefficient functions may have discontinuity at finite interior point. Naturally, eigenfunctions of this problem may have discontinuity at the finite inner point of the considered interval. The problems with transmission conditions has become an important area of research in recent years because of the needs of modern technology, engineering and physics. Many of the mathematical problems encountered in the study of boundary-value-transmission problem cannot be treated with the usual techniques within the standard framework of boundary value problem (see [**?**]). Note that some special cases of this problem arise after an application of the method of separation of variables to a varied assortment of physical problems. For example, some boundary value problems with transmission conditions arise in heat and mass transfer problems [1], in vibrating string problems when the string loaded additionally with point masses [7], in diffraction problems [10]. Also some problems with transmission conditions which arise in mechanics (thermal conduction problems for a thin laminated plate) were studied in [9].

2. The fundamental solutions and characteristic function

With a view to constructing the characteristic function $\omega(\lambda)$ we shall define two fundamental solution

$$\phi(x,\lambda)=\begin{cases}\phi_1(x,\lambda), x\in[a,\xi_1)\\ \phi_2(x,\lambda), x\in(\xi_1,\xi_2)\\ \phi_3(x,\lambda), x\in(\xi_2,\xi_3)\\ \quad ...\\ \phi_{r+1}(x,\lambda), x\in(\xi_r,b]\end{cases} \text{ and } \chi(x,\lambda)=\begin{cases}\chi_1(x,\lambda), x\in[a,\xi_1)\\ \chi_2(x,\lambda), x\in(\xi_1,\xi_2)\\ \chi_3(x,\lambda), x\in(\xi_2,\xi_3)\\ ...\\ \chi_{r+1}(x,\lambda), x\in(\xi_r,b]\end{cases}$$

by special procedure. Let $\phi_1(x,\lambda)$ and $\chi_{r+1}(x,\lambda)$ be solutions of the equation (1) on (a,ξ_1) and (ξ_n,b) satisfying initial conditions

$$u(a,\lambda)=\alpha_2-\lambda\alpha_4,\ u'(a,\lambda)=\alpha_1-\lambda a_3 \tag{6}$$

$$u(b,\lambda)=\beta_2+\lambda\beta_4,\ u'(b,\lambda)=\beta_1+\lambda\beta_3 \tag{7}$$

respectively. In terms of these solution we shall define the other solutions $\phi_i(x,\lambda)$, and $\chi_i(x,\lambda)$ by initial conditions

$$\phi_{i+1}(\xi_i+,\lambda)=\frac{1}{\Delta_{i12}}(\Delta_{i23}\phi_i(\xi_i-,\lambda)+\Delta_{i24}\frac{\partial\phi_i(\xi_i-,\lambda)}{\partial x}) \tag{8}$$

$$\frac{\partial\phi_{i+1}(\xi_i+,\lambda)}{\partial x}=\frac{-1}{\Delta_{i12}}(\Delta_{i13}\phi_i(\xi_i-,\lambda)+\Delta_{i14}\frac{\partial\phi_i(\xi_i-,\lambda)}{\partial x}) \tag{9}$$

and

$$\chi_i(\xi_i-,\lambda)=\frac{-1}{\Delta_{i34}}(\Delta_{i14}\chi_{i+1}(\xi_i+,\lambda)+\Delta_{i24}\frac{\partial\chi_{i+1}(\xi_i+,\lambda)}{\partial x}) \tag{10}$$

$$\frac{\partial\chi_i(\xi_i-,\lambda)}{\partial x})=\frac{1}{\Delta_{i34}}(\Delta_{i13}\chi_{i+1}(\xi_i+,\lambda)+\Delta_{i23}\frac{\partial\chi_{i+1}(\xi_i+,\lambda)}{\partial x}) \tag{11}$$

respectively, where Δ_{ijk} $(1\leq j<k\leq 4)$ denotes the determinant of the j-th and k-th columns of the matrix

$$\begin{bmatrix}\alpha_{i1}^{+} & \alpha_{i0}^{+} & \alpha_{i1}^{-} & \alpha_{i0}^{-}\\ \beta_{i1}^{+} & \beta_{i0}^{+} & \beta_{i1}^{-} & \beta_{i0}^{-}\end{bmatrix}$$

where $i=1,2,...r$. Everywhere in below we shall assume that $\Delta_{ijk}>0$ for all i,j,k. The existence and uniqueness of these solutions are follows from well-known theorem of ordinary differential equation theory. Moreover by applying the method of [4] we can prove that each of these solutions are entire functions of parameter $\lambda \in \mathbb{C}$ for each fixed x. Taking into account (8)-(11) and the fact that the Wronskians $\omega_i(\lambda) := W[\phi_i(x,\lambda), \chi_i(x,\lambda)]$ (i=1,2,...r+1) are independent of variable x we have

$$\begin{aligned}
\omega_{i+1}(\lambda) &= \phi_i(\xi_i+,\lambda)\frac{\partial\chi_i(\xi_i+,\lambda)}{\partial x} - \frac{\partial\phi_i(\xi_i+,\lambda)}{\partial x}\chi_i(\xi_i+,\lambda) \\
&= \frac{\Delta_{i34}}{\Delta_{i12}}(\phi_i(\xi_i-,\lambda)\frac{\partial\chi_i(\xi_i-,\lambda)}{\partial x} - \frac{\partial\phi_i(\xi_i-,\lambda)}{\partial x}\chi_i(\xi_i-,\lambda)) \\
&= \frac{\Delta_{i34}}{\Delta_{i12}}\omega_i(\lambda) = \prod_{j=1}^{i}\frac{\Delta_{j34}}{\Delta_{j12}}\omega_1(\lambda)\ (i=1,2,...r).
\end{aligned}$$

It is convenient to define the characteristic function $\omega(\lambda)$ for our problem $(1)-(3)$ as

$$\omega(\lambda) := \omega_1(\lambda) = \prod_{j=1}^{i}\frac{\Delta_{j12}}{\Delta_{i34}}\omega_{i+1}(\lambda)\ (i=1,2,...r).$$

Obviously, $\omega(\lambda)$ is an entire function. By applying the technique of [3] we can prove that there are infinitely many eigenvalues $\lambda_n,\ n=1,2,...$ of the problem $(1)-(5)$ which are coincide with the zeros of characteristic function $\omega(\lambda)$.

3. Some results for eigenvalues

Theorem 1. *The eigenvalues of the problem* (1)-(5) *are consist of the zeros of the function* $w(\lambda)$.

Theorem 2. *The eigenvalues of the problem* (1)-(5) *are are analytically simple.*

Theorem 3. *The eigenvalues of the problem* (1)-(5) *are also geometrically simple.*

Theorem 4. *The eigenvalues of the problem* (1)-(5) *are bounded below, and they are countably infinite and can cluster only at* ∞.

Now by modifying the standard method we prove that all eigenvalues of the problem $(1)-(5)$ are real.

Theorem 5. *The eigenvalues of the boundary value transmission problem* $(1)-(4)$ *are real.*

4. Asymptotic formulas for ,,basic" solutions

Below, for shorting we shall use also notations; $\phi_i(x,\lambda) := \phi_{i\lambda}(x),\ \chi_i(x,\lambda) := \chi_{i\lambda}(x)$ $(i = 1,2,...r+1)$. We can prove that the next integral and integro-differential equations are hold for $k=0$ and $k=1$.

$$\begin{aligned}
\frac{d^k}{dx^k}\phi_{1\lambda}(x) &= (\alpha_2 - s^2\alpha_4)\frac{d^k}{dx^k}\cos\left[\frac{s\,(x-a)}{\rho_1}\right] + \frac{\rho_1(\alpha_1 - s^2\alpha_3)}{s}\frac{d^k}{dx^k}\sin\left[\frac{s\,(x-a)}{\rho_1}\right] \\
&+ \frac{1}{\rho_1 s}\int\limits_a^x \frac{d^k}{dx^k}\sin\left[\frac{s\,(x-y)}{\rho_1}\right] q(y)\phi_{1\lambda}(y)dy, \qquad (12) \\
\frac{d^k}{dx^k}\phi_{(i+1)\lambda}(x) &= \frac{1}{\Delta_{i12}}(\Delta_{i23}\phi_{i\lambda}(\xi_i) + \Delta_{i24}\phi'_{i\lambda}(\xi_i))\frac{d^k}{dx^k}\cos\left[\frac{s(x-\xi_i)}{\rho_{i+1}}\right] \\
&- \frac{\rho_{i+1}}{s\Delta_{i12}}(\Delta_{i13}\phi_{i\lambda}(\xi_i) + \Delta_{i14}\phi'_{i\lambda}(\xi_i))\frac{d^k}{dx^k}\sin\left[\frac{s(x-\xi_i)}{\rho_{i+1}}\right] \\
&+ \frac{1}{\rho_{i+1} s}\int\limits_{\xi_i}^x \frac{d^k}{dx^k}\sin\left[\frac{s\,(x-y)}{\rho_{i+1}}\right] q(y)\phi_{(i+1)\lambda}(y)dy \qquad (13)
\end{aligned}$$

for x $\in (a,\xi_1)$ and $x \in (\xi_i,\xi_{i+1}),\ i = 1,2,...r$ respectively.

$$\begin{aligned}
\frac{d^k}{dx^k}\chi_{(r+1)\lambda}(x) &= (\beta_2 + s^2\beta_4)\frac{d^k}{dx^k}\cos\left[\frac{s\,(x-b)}{\rho_{r+1}}\right] + \frac{\rho_{r+1}}{s}(\beta_2 + s^2\beta_3)\frac{d^k}{dx^k}\sin\left[\frac{s\,(x-b)}{\rho_{r+1}}\right] \\
&+ \frac{1}{\rho_{r+1} s}\int\limits_x^b \frac{d^k}{dx^k}\sin\left[\frac{s\,(x-y)}{\rho_{r+1}}\right] q(y)\chi_{(r+1)\lambda}(y)dy \qquad (14) \\
\frac{d^k}{dx^k}\chi_{i\lambda}(x) &= -\frac{1}{\Delta_{i34}}(\Delta_{i14}\chi_{(i+1)\lambda}(\xi_i) + \Delta_{i24}\chi'_{(i+1)\lambda}(\xi_i))\frac{d^k}{dx^k}\cos\left[\frac{s(x-\xi_i)}{\rho_i}\right] \\
&- \frac{\rho_i}{s\Delta_{i34}}(\Delta_{i13}\chi_{(i+1)\lambda}(\xi_i) + \Delta_{i23}\chi'_{(i+1)\lambda}(\xi_i))\frac{d^k}{dx^k}\sin\left[\frac{s(x-\xi_i)}{\rho_i}\right] \\
&+ \frac{1}{\rho_i s}\int\limits_x^{\xi_i} \frac{d^k}{dx^k}\sin\left[\frac{s\,(x-y)}{\rho_i}\right] q(y)\chi_{i\lambda}(y)dy \qquad (15)
\end{aligned}$$

for $x \in (\xi_n, b)$ and (ξ_i,ξ_{i+1}) (i=1,2,...r) respectively.

Theorem 6. *Let $Im\mu = t$. Then if $\alpha_4 \neq 0$*

$$\frac{d^k}{dx^k}\phi_{1\lambda}(x) = -\alpha_4 s^2 \frac{d^k}{dx^k}\cos\left[\frac{s\,(x-a)}{\rho_1}\right] + O\left(|s|^{k+1}\, e^{|t|\frac{(x-a)}{\rho_1}}\right) \tag{16}$$

$$\begin{aligned} \frac{d^k}{dx^k}\phi_{(i+1)\lambda}(x) &= (-1)^{i+1}\alpha_4 s^{i+2}\left(\prod_{j=1}^{i}\frac{\Delta_{j24}}{\Delta_{j12}\rho_j}\sin\left[\frac{s\,(\xi_j-\xi_{j-1})}{\rho_j}\right]\right)\frac{d^k}{dx^k}\cos\left[\frac{s\,(x-\xi_i)}{\rho_{i+1}}\right] \\ &\quad +O\left(|s|^{i+1}e^{|t|((\sum\limits_{j=1}^{i}\frac{(\xi_j-\xi_{j-1})}{\rho_j})+\frac{(x-\xi_i)}{\rho_{i+1}})}\right), \quad i=1,2,...r \end{aligned} \tag{17}$$

as $|s| \to \infty$, while if $\alpha_4 = 0$

$$\frac{d^k}{dx^k}\phi_{1\lambda}(x) = -\alpha_3 s \frac{d^k}{dx^k}\sin\left[\frac{s\,(x-a)}{\rho_1}\right] + O\left(|s|^{k}\, e^{|t|\frac{(x-a)}{\rho_1}}\right) \tag{18}$$

$$\begin{aligned} \frac{d^k}{dx^k}\phi_{2\lambda}(x) &= -\alpha_3 s^2\frac{\Delta_{124}}{\Delta_{112}}\frac{d^k}{dx^k}\cos\left[\frac{s\,(\xi_1-a)}{\rho_1}\right]\cos\left[\frac{s\,(x-\xi_1)}{\rho_2}\right] \\ &+ O\left(|s|^{k}\, e^{|t|(\frac{(\xi_1-a)}{\rho_1}+\frac{(x-\xi_1)}{\rho_2})}\right) \end{aligned} \tag{19}$$

$$\begin{aligned} \frac{d^k}{dx^k}\phi_{(i+1)\lambda}(x) &= (-1)^{i}\alpha_3 s^{i+1}\cos\left[\frac{s\,(\xi_1-a)}{\rho_1}\right]\left(\prod_{j=2}^{i}\frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j}\sin\left[\frac{s\,(\xi_j-\xi_{j-1})}{\rho_j}\right]\right) \\ &\times \frac{d^k}{dx^k}\cos\left[\frac{s\,(x-\xi_i)}{\rho_{i+1}}\right] + O\left(|s|^{i}e^{|t|((\sum\limits_{j=1}^{i}\frac{(\xi_j-\xi_{j-1})}{\rho_j})+\frac{(x-\xi_j)}{\rho_{j+1}})}\right) \end{aligned} \tag{20}$$

$i = 2, ...r$ as $|s| \to \infty$ ($k = 0, 1$). Each of this asymptotic equalities hold uniformly for x.

Theorem 7. *Let $Ims = t$. Then if $\beta_4 \neq 0$*

$$\frac{d^k}{dx^k}\chi_{(r+1)\lambda}(x) = \beta_4 s^2 \frac{d^k}{dx^k}\cos\left[\frac{s\,(b-x)}{\rho_{r+1}}\right] + O\left(|s|^{k+1}\, e^{|t|\frac{(b-x)}{\rho_{r+1}}}\right) \tag{21}$$

$$\begin{aligned} \frac{d^k}{dx^k}\chi_{(r-i)\lambda}(x) &= (-1)^{i+1}\beta_4 s^{i+3}\left(\prod_{j=0}^{i}\frac{\Delta_{(r-j)24}}{\Delta_{(r-j)34}\rho_{r+1-j}}\sin\left[\frac{s\,(\xi_{r+1-j}-\xi_{r-j})}{\rho_{r+1-j}}\right]\right) \\ &\times \frac{d^k}{dx^k}\cos\left[\frac{s\,(x-\xi_{r-i})}{\rho_{r-i}}\right] + O\left(|s|^{i+2}e^{|t|((\sum\limits_{j=0}^{i}\frac{(\xi_{r+1-j}-\xi_{r-j})}{\rho_{r+1-j}})+\frac{(x-\xi_{r-j})}{\rho_{r-j}})}\right), \end{aligned} \tag{22}$$

as $|s| \to \infty$*, while if* $\beta_4 = 0$

$$\begin{aligned}
\frac{d^k}{dx^k}\chi_{(r+1)\lambda}(x) &= -\beta_3\rho_{r+1}s\frac{d^k}{dx^k}\sin\left[\frac{s\,(b-x)}{\rho_{r+1}}\right] + O\left(|s|^{k+1}e^{|t|\frac{(b-x)}{\rho_{r+1}}}\right) &(23)\\
\frac{d^k}{dx^k}\chi_{r\lambda}(x) &= -\beta_3 s^2\frac{\Delta_{r24}}{\Delta_{r34}}\cos\left[\frac{s\,(b-\xi_r)}{\rho_{r+1}}\right]\cos\left[\frac{s\,(x-\xi_r)}{\rho_r}\right]\\
&+ O\left(|s|^{k+1}e^{|t|(\frac{(b-\xi_r)}{\rho_{r+1}}+\frac{(x-\xi_r)}{\rho_r})}\right) &(24)\\
\frac{d^k}{dx^k}\chi_{(r-i)\lambda}(x) &= (-1)^{i+1}\beta_3 s^{i+2}\cos\left[\frac{s\,(b-\xi_r)}{\rho_{r+1}}\right]\frac{d^k}{dx^k}\cos\left[\frac{s\,(x-\xi_{r-i})}{\rho_{r-i}}\right]\\
&\times \left(\prod_{j=0}^{i}\frac{\Delta_{(r-j)24}}{\Delta_{(r-j)34}\rho_{r-j}}\sin\left[\frac{s\,(\xi_{r-1-j}-\xi_{r-j})}{\rho_{r-j}}\right]\right)\\
&+ O\left(|s|^{i+1}e^{|t|((\sum\limits_{j=0}^{i}\frac{(\xi_{r+1-j}-\xi_{r-j})}{\rho_{r+1-j}})+\frac{(x-\xi_{r-j})}{\rho_{r-j}})}\right) &(25)
\end{aligned}$$

as $|s| \to \infty$ $i = 1, ...r-1$ $(k = 0, 1)$.*Each of this asymptotic equalities hold uniformly for* x.

5. Asymptotic behaviour of eigenvalues and eigenfunctions

Since the Wronskians of $\phi_\lambda(x)$ and $\chi_\lambda(x)$ are independent of x in each $\Omega_i (i = 0, 1, ..., r+1)$, in particular, by putting $x = b$ we have

$$\begin{aligned}
\omega(\lambda) &= \prod_{j=1}^{r}\frac{\Delta_{j12}}{\Delta_{j34}}\omega_{r+1}(\lambda)|_{x=b} = \prod_{j=1}^{r}\frac{\Delta_{j12}}{\Delta_{j34}}\omega(\phi_{r+1}(b,\lambda),\chi_{r+1}(b,\lambda))\\
&= \prod_{j=1}^{r}\frac{\Delta_{j12}}{\Delta_{j34}}\{(\beta_1+\lambda\beta_3)\phi_{(n+1)}(b,\lambda)+(\beta_2+\lambda\beta_4)\phi'_{(n+1)}(b,\lambda)\}. &(26)
\end{aligned}$$

Let $Im\mu = t$. By substituting (21) and (25) in (26) we obtain the following asymptotic representations

(i) If $\alpha_4 \neq 0$ and $\beta_4 \neq 0$, then

$$w(\lambda) = -\alpha_4\beta_4 s^{r+5}\left(\prod_{j=1}^{r+1}\frac{1}{\rho_j}\sin\left[\frac{s\,(\xi_{j-1}-\xi_j)}{\rho_j}\right]\right) + O\left(|s|^{r+4}e^{|t|(\sum\limits_{j=1}^{r+1}\frac{(\xi_j-\xi_{j-1})}{\rho_j})}\right) \tag{27}$$

(ii) If $\alpha_4 \neq 0$ and $\beta_4 = 0$, then

$$\begin{aligned} w(\lambda) &= -\alpha_4\beta_3 s^{r+4}\cos\left[\frac{s\left(b-\xi_r\right)}{\rho_{r+1}}\right]\left(\prod_{j=1}^{r}\frac{1}{\rho_j}\sin\left[\frac{s\left(\xi_{j-1}-\xi_j\right)}{\rho_j}\right]\right) \\ &+ O\left(|s|^{r+3}e^{|t|(\sum\limits_{j=1}^{r+1}\frac{(\xi_j-\xi_{j-1})}{\rho_j})}\right) \end{aligned} \tag{28}$$

(iii) If $\alpha_4 = 0$ and $\beta_4 \neq 0$, then

$$\begin{aligned} w(\lambda) &= -\alpha_3\beta_4 s^{r+4}\frac{\Delta_{112}}{\Delta_{134}}\cos\left[\frac{s\left(\xi_1-a\right)}{\rho_1}\right]\left(\prod_{j=2}^{r+1}\frac{1}{\rho_j}\sin\left[\frac{s\left(\xi_{j-1}-\xi_j\right)}{\rho_j}\right]\right) \\ &+ O\left(|s|^{r+3}e^{|t|(\sum\limits_{j=1}^{r+1}\frac{(\xi_j-\xi_{j-1})}{\rho_j})}\right) \end{aligned} \tag{29}$$

(iv) If $\alpha_4 = 0$ and $\beta_4 = 0$, then

$$\begin{aligned} w(\lambda) &= -\alpha_3\beta_3 s^{r+3}\frac{\Delta_{112}}{\Delta_{134}}\cos\left[\frac{s\left(\xi_1-a\right)}{\rho_1}\right]\cos\left[\frac{s\left(b-\xi_r\right)}{\rho_{r+1}}\right] \\ &\times \left(\prod_{j=2}^{r}\frac{1}{\rho_{j-1}}\sin\left[\frac{s\left(\xi_{j-1}-\xi_j\right)}{\rho_j}\right]\right) + O\left(|s|^{r+2}e^{|t|(\sum\limits_{j=1}^{r+1}\frac{(\xi_j-\xi_{j-1})}{\rho_j})}\right) \end{aligned} \tag{30}$$

Now we are ready to derived the needed asymptotic formulas for eigenvalues and eigenfunctions.

Theorem 8. *The boundary-value-transmission problem* (1)-(5) *has an precisely numerable many real eigenvalues, whose behavior may be expressed by* $r+1$ *sequence* $\{s_n^j\}$ $(j=1,2,..r+1)$ *with following asymptotic as* $n \to \infty$
(i) *If* $\alpha_4 \neq 0$ *and* $\beta_4 \neq 0$, *then*

$$s_n^{(j)} = (\frac{n}{2}-1)\frac{\rho_j\pi}{(\xi_{j-1}-\xi_j)} + O\left(\frac{1}{n}\right), \quad (j=1,2,...r+1) \tag{31}$$

(ii) *If* $\alpha_4 \neq 0$ *and* $\beta_4 = 0$, *then*

$$s_n^{(r+1)} = (n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_r)}+O\left(\frac{1}{n}\right),\ s_n^{(j)} = \frac{(n-1)\rho_j\pi}{2(\xi_{j-1}-\xi_j)}+O\left(\frac{1}{n}\right),\ (j=1,...r), \tag{32}$$

(iii) *If $\alpha_4 = 0$ and $\beta_4 \neq 0$, then*

$$s_n^{(1)} = (n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1 - a)}+O\left(\frac{1}{n}\right), \ s_n^{(j)} = \frac{(n-1)\rho_j\pi}{2(\xi_{j-1}-\xi_j)}+O\left(\frac{1}{n}\right), \ \ (j = 2, ...r+1), \tag{33}$$

(iv) *If $\alpha_4 = 0$ and $\beta_4 = 0$, then*

$$\begin{aligned} s_n^{(1)} &= (n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1 - a)}+O\left(\frac{1}{n}\right), \ s_n^{(r+1)} = (n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_r)}+O\left(\frac{1}{n}\right), \\ s_n^{(j)} &= \frac{n\rho_j\pi}{2(\xi_{j-1}-\xi_j)}+O\left(\frac{1}{n}\right), \ \ (j = 2, ...r), \end{aligned} \tag{34}$$

Proof. Let $\alpha_4 \neq 0$ and $\beta_4 \neq 0$. By applying the well-known Rouche Theorem which asserts that if $f(z)$ and $g(z)$ are analytic inside and on a closed contour Γ, and $|g(z)| < |f(z)|$ on Γ then $f(z)$ and$f(z) + g(z)$ have the same number zeros inside Γ provided that the zeros are counted with multiplicity on a sufficiently large contour, it follows that $w(\lambda)$ has the same number of zeros inside the suitable contour as the leading term $w_0(\lambda) = -\alpha_4\beta_4 s^{r+5}\left(\prod\limits_{j=1}^{r+1}\frac{1}{\rho_j}\sin\left[\frac{s(\xi_{j-1}-\xi_j)}{\rho_j}\right]\right)$ in (27). Hence, if $\lambda_0 < \lambda_1 < \lambda_2 ...$ are the zeros of $w(\lambda)$ and λ_n, we have the needed asymptotic formulas (31). Other cases can be proved similarly.

Using this asymptotic expressions of eigenvalues we can obtain the corresponding asymptotic expressions for eigenfunctions of the problem (1)-(4). Recalling that $\phi_{\lambda_n}(x)$ is an eigenfunction according to the eigenvalue λ_n, and by putting (31) in the (16)-(17) for $k = 0, 1$ and denoting the corresponding eigenfunction as $\phi_n^{(k)}(x)$ $k = 1, 2, ...r+1$ we get the following cases If $\alpha_4 \neq 0$ and $\beta_4 \neq 0$, then

$$\phi_n^{(k)}(x) = \begin{cases} -\alpha_4\left[(\frac{n}{2}-1)\frac{\rho_k\pi}{(\xi_{k-1}-\xi_k)}\right]^2\cos\left[(\frac{n}{2}-1)\frac{\rho_k\pi}{(\xi_{k-1}-\xi_k)}\frac{(x-a)}{\rho_1}\right]+O(n), \ \ x\in(a,\xi_1) \\ (-1)^{i+1}\alpha_4\left[(\frac{n}{2}-1)\frac{\rho_k\pi}{(\xi_{k-1}-\xi_k)}\right]^{i+2}\left(\prod_{j=1}^{i}\frac{\Delta_{j24}}{\Delta_{j12}\rho_j}\sin\left[(\frac{n}{2}-1)\frac{\rho_k\pi(\xi_j-\xi_{j-1})}{(\xi_{k-1}-\xi_k)\rho_j}\right]\right) \\ \times\cos\left[(\frac{n}{2}-1)\frac{\rho_k\pi(x-\xi_i)}{(\xi_{k-1}-\xi_k)\rho_{i+1}}\right]+O(n^{i+1}), \ \ x\in(\xi_i,\xi_{i+1}), \ i = 1, 2, ...r \end{cases}$$

where k=1,2,...r+1. If $\alpha_4 \neq 0$ and $\beta_4 = 0$

$$\phi_n^{(r+1)}(x) = \begin{cases} -\alpha_4 \left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_n)}\right]^2 \cos\left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_n)}\frac{(x-a)}{\rho_1}\right] + O(n), \quad x \in (a,\xi_1) \\ (-1)^{i+1}\alpha_4 \left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_n)}\right]^{i+2} \left(\prod_{j=1}^{i} \frac{\Delta_{j24}}{\Delta_{j12}\rho_j} \sin\left[(\frac{1}{2}+n)\frac{\rho_{r+1}\pi(\xi_j-\xi_{j-1})}{(b-\xi_n)\rho_j}\right]\right) \\ \times \cos\left[(\frac{1}{2}+n)\frac{\rho_{r+1}\pi(x-\xi_i)}{(b-\xi_n)\rho_{i+1}}\right] + O(n^{i+1}), \ x \in (\xi_i,\xi_{i+1}) \ \ i = 1,2,...r \end{cases}$$

and

$$\phi_n^{(k)}(x) = \begin{cases} -\alpha_4 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^2 \cos\left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\frac{(x-a)}{\rho_1}\right] + O(n), \quad x \in (a,\xi_1) \\ (-1)^{i+1}\alpha_4 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^{i+2} \left(\prod_{j=1}^{i} \frac{\Delta_{j24}}{\Delta_{j12}\rho_j} \sin\left[\frac{(n-1)\rho_k\pi(\xi_j-\xi_{j-1})}{2(\xi_{k-1}-\xi_k)\rho_j}\right]\right) \\ \times \cos\left[\frac{(n-1)\rho_k\pi(x-\xi_i)}{2(\xi_{k-1}-\xi_k)\rho_{i+1}}\right] + O(n^{i+1}), \quad i = 1,2,...r \text{ for } x \in (\xi_i,\xi_{i+1}) \end{cases}$$

where k=1,2,...r. If $\alpha_4 = 0$ and $\beta_4 \neq 0$, then

$$\phi_n^{(1)}(x) = \begin{cases} -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right] \sin\left[(n+\frac{1}{2})\frac{\rho_1\pi(x-a)}{(\xi_1-a)\rho_1}\right] + O(1), \quad x \in (a,\xi_1) \\ -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right]^2 \frac{\Delta_{124}}{\Delta_{112}} \cos\left[(n+\frac{1}{2})\pi\right] \cos\left[\frac{(n+\frac{1}{2})\rho_1\pi(x-\xi_1)}{(\xi_1-a)\rho_2}\right] + O(n) \\ \text{for } x \in (\xi_1,\xi_2) \\ (-1)^i\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right]^{i+1} \cos\left[(n+\frac{1}{2})\pi\right] \cos\left[(n+\frac{1}{2})\frac{\rho_1\pi(x-\xi_i)}{(\xi_1-a)\rho_{i+1}}\right] \\ \times \left(\prod_{j=2}^{i} \frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j} \sin\left[(n+\frac{1}{2})\frac{\rho_1\pi(\xi_j-\xi_{j-1})}{(\xi_1-a)\rho_j}\right]\right) + O(n^i) \ x \in (\xi_i,\xi_{i+1}) \\ i = 2,...r \end{cases}$$

and

$$\phi_n^{(k)}(x) = \begin{cases} -\alpha_3 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right] \sin\left[\frac{(n-1)\rho_k\pi(x-a)}{2(\xi_{k-1}-\xi_k)\rho_1}\right] + O(1), \quad x \in (a,\xi_1) \\ -\alpha_3 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^2 \frac{\Delta_{124}}{\Delta_{112}} \cos\left[\frac{(n-1)(\xi_1-a)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_1}\right] \cos\left[\frac{(n-1)(x-\xi_1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_2}\right] + O(n) \\ \text{for } x \in (\xi_1,\xi_2) \\ (-1)^i\alpha_3 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^{i+1} \cos\left[\frac{(n-1)(\xi_1-a)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_1}\right] \cos\left[\frac{(n-1)(x-\xi_i)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_{i+1}}\right] \\ \times \left(\prod_{j=2}^{i} \frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j} \sin\left[\frac{(n-1)(\xi_j-\xi_{j-1})\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_j}\right]\right) + O(n^i), \quad x \in (\xi_i,\xi_{i+1}) \\ i = 2,...r \end{cases}$$

where k=2,...r+1. If $\alpha_4 = 0$ and $\beta_4 = 0$, then

$$\phi_n^{(1)}(x) = \begin{cases} -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right] \sin\left[(n+\frac{1}{2})\frac{\rho_1\pi(x-a)}{(\xi_1-a)\rho_1}\right] + O(1), \quad x \in (a,\xi_1) \\ -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right]^2 \frac{\Delta_{124}}{\Delta_{112}} \cos\left[(n+\frac{1}{2})\pi\right] \cos\left[\frac{(n+\frac{1}{2})\rho_1\pi(x-\xi_1)}{(\xi_1-a)\rho_2}\right] + O(n) \\ \text{for } x \in (\xi_1,\xi_2) \\ (-1)^i\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_1\pi}{(\xi_1-a)}\right]^{i+1} \cos\left[(n+\frac{1}{2})\pi\right] \cos\left[(n+\frac{1}{2})\frac{\rho_1\pi(x-\xi_i)}{(\xi_1-a)\rho_{i+1}}\right] \\ \times\left(\prod_{j=2}^{i} \frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j} \sin\left[(n+\frac{1}{2})\frac{\rho_1\pi(\xi_j-\xi_{j-1})}{(\xi_1-a)\rho_j}\right]\right) + O(n^i), \; x \in (\xi_i,\xi_{i+1}) \\ i = 2,...r \end{cases}$$

$$\phi_n^{(r+1)}(x) = \begin{cases} -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_n)}\right] \sin\left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi(x-a)}{(b-\xi_n)\rho_1}\right] + O(1), \quad x \in (a,\xi_1) \\ -\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(b-\xi_n)}\right]^2 \frac{\Delta_{124}}{\Delta_{112}} \cos\left[\frac{(n+\frac{1}{2})\rho_{r+1}\pi(\xi_1-a)}{(b-\xi_n)\rho_1}\right] \cos\left[\frac{(n+\frac{1}{2})\rho_{r+1}\pi(x-\xi_1)}{(b-\xi_n)\rho_2}\right] \\ +O(n) \text{ for } x \in (\xi_1,\xi_2) \\ (-1)^i\alpha_3 \left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi}{(\xi_1-a)}\right]^{i+1} \cos\left[\frac{(n+\frac{1}{2})\rho_{r+1}\pi(\xi_1-a)}{(b-\xi_n)\rho_1}\right] \cos\left[(n+\frac{1}{2})\frac{\rho_{r+1}\pi(x-\xi_i)}{(b-\xi_n)\rho_{i+1}}\right] \\ \times\left(\prod_{j=2}^{i} \frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j} \sin\left[\frac{(n+\frac{1}{2})\rho_{r+1}\pi(\xi_j-\xi_{j-1})}{(b-\xi_n)\rho_j}\right]\right) + O(n^i), \quad x \in (\xi_i,\xi_{i+1}) \\ i = 2,...r \end{cases}$$

and

$$\phi_n^{(k)}(x) = \begin{cases} -\alpha_3 \left[\frac{(n-1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right] \sin\left[\frac{(n-1)\rho_k\pi(x-a)}{2(\xi_{k-1}-\xi_k)\rho_1}\right] + O(1), \quad x \in (a,\xi_1) \\ -\alpha_3 \left[\frac{n\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^2 \frac{\Delta_{124}}{\Delta_{112}} \cos\left[\frac{n(\xi_1-a)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_1}\right] \cos\left[\frac{n(x-\xi_1)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_2}\right] + O(n) \\ (-1)^i\alpha_3 \left[\frac{n\rho_k\pi}{2(\xi_{k-1}-\xi_k)}\right]^{i+1} \cos\left[\frac{n(\xi_1-a)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_1}\right] \cos\left[\frac{n(x-\xi_i)\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_{i+1}}\right] \\ \times\left(\prod_{j=2}^{i} \frac{\Delta_{(j-1)24}}{\Delta_{(j-1)12}\rho_j} \sin\left[\frac{n(\xi_j-\xi_{j-1})\rho_k\pi}{2(\xi_{k-1}-\xi_k)\rho_j}\right]\right) + O(n^i), \quad x \in (\xi_i,\xi_{i+1}) \\ i = 2,...r \end{cases}$$

where k=2,...r+1. All this asymptotic approximations are hold uniformly for x.

References

[1] A. V. Likov and Yu. A. Mikhailov, *The theory of Heat and Mass Transfer*, Qosenergaizdat, 1963(Russian).

[2] O. Sh. Mukhtarov and H. Demir, *Coersiveness of the discontinuous initial- boundary value problem for parabolic equations*, Israel J. Math., Vol. 114(1999), Pages 239-252.

[3] M. Kadakal and O. Sh. Mukhtarov *SturmLiouville problems with discontinuities at two points*, Computers and Mathematics with Applications 54(2007) 1367-1379

[4] F. S. Muhtarov and K. Aydemir*Distributions of eigenvalues for Sturm-Liouville problem under jump conditions*, Journal of New Results in Science 1(2012) 81-89.

[5] O. Sh. Mukhtarov and S. Yakubov, *Problems for ordinary differential equations with transmission conditions*, Appl. Anal.,81(2002),1033-1064.

[6] M. L. Rasulov, *Methods of Contour Integration*, North-Holland Publishing Company, Amsterdam, 1967.

[7] A. N. Tikhonov and A. A. Samarskii, *Equations of Mathematical Physics,* Oxford and New York, Pergamon, 1963.

[8] E. C. Titchmarsh, *Eigenfunctions Expansion Associated with Second Order Differential Equations I*, second edn. Oxford Univ. Press, London, 1962.

[9] I. Titeux and Ya. Yakubov, *Completeness of root functions for thermal conduction in a strip with piecewise continuous coefficients*, Math. Models Methods Appl. Sc., 7(7), (1997), 1035-1050.

[10] N. N. Voitovich , B. Z. Katsenelbaum and A. N. Sivov , *Generalized Method of Eigen-vibration in the theory of Diffraction* , Nakua, Mockow, 1997 (Russian).